



Focus Group Fertiliser efficiency in horticulture

Mini-paper - Need for system approach to increase nutrient use efficiency in horticulture

Janjo de Haan¹, Stefaan De Neve², Eligio Malusa³, Fernando Andres Torresano-Sanchez⁴, Zoltàn Hadju⁵

¹ Applied Plant Research – Wageningen UR, The Netherlands

² University of Ghent, Belgium

³ Research Institute of Horticulture, Poland and Agricultural Research Council, Italy

⁴ Nutricrop, Spain

⁵ SOLTUB Kft., Hungary

Introduction

Higher nutrient efficiency means that a larger share of the fertiliser input is utilised by the crop and is ending in the output, products harvested from the field. There is a large need to increase the nutrient efficiency of horticultural crops. Firstly, to reduce emissions of ammonia and nitrous oxides related to fertiliser use. Secondly to improve ground water and surface water quality as required by the EU Nitrates Directive (91/676/EEC) and the EU Water Framework Directive (2000/60/EC). An increase in nutrient use efficiency is needed because inputs from fertilisers are becoming scarce: e.g. phosphate rock but also energy to produce nitrogen fertilisers (IFADATA, 2014). Finally an increase in fertiliser efficiency is also important to reduce production costs and increase profits for farmers.

Various measures are available to improve nutrient efficiency. However, simple solutions are often lacking as different aspects have to be taken into account when starting to select the appropriate measure(s): the objectives for which nutrient use efficiency has to be increased, the effect of the different measures under local biophysical conditions, the feasibility and the costs and benefits for farmers. Very often, trade-offs between objectives need to be considered in selecting the measures. Besides, measures can have unwanted direct or indirect side effects outside the scope of the objectives. A direct side effect is pollution swapping, for instance the reduction of NH₃ emissions by injection of animal manures may increase N₂O losses and leaching. An example of an indirect side effect is the application of drip-irrigation, which makes mechanical weed control more difficult, eventually increasing the use and emission of herbicides. Finally, it is important to note that the main objectives of farmers are producing good quality crops and making a profit and not improving nutrient efficiency. The costs of fertilisers are yet not a major obstacle and the fulfilment of EU requirements is mainly administrative and not leading to change of behaviour. Therefore, farmers are not often motivated to reduce the use of fertilisers.

A system approach is needed to make a coherent integrated solution that reaches the objectives of both farmers and society as best as possible. This system approach should focus on the farm level as the farmer is responsible to select the measures that will be applied in the field. In this paper, we describe how the prototyping approach can be used to improve effectively nutrient use efficiency.



The approach is extensive. Therefore, we discuss various examples that uses parts of the approach. To conclude we provide some recommendations on how system approaches can be used more extensively to improve overall sustainability of horticulture production.

Development of farming systems (prototyping)

For the development of farming systems for vegetable crops, a standardised methodology called "prototyping" has been developed. The methodology is a combined research and development effort beginning with a profile of agronomic, environmental and economic demands (objectives) for a more sustainable, future-oriented farming. This exercise ends with tested, ready-to-use prototypes, designed for widespread use. The prototyping methodology was examined for arable farming in a four-year European Union Concerted Action (Vereijken, 1999) and further developed for vegetable farming during the EU-VEGINECO-project (de Haan & Garcia Diaz, 2002; de Haan, 2002).

The methodology consists of four phases: 1) analysis and diagnosis, 2) design, 3) testing and improving and 4) dissemination (Figure 1). The first phase starts with a regionally based analysis of the prevailing agricultural systems resulting in a diagnosis of the situation in terms of sectorial statistics, farm structure, agro-ecological state-of-the-art, ecological–environmental impact, the socio-economic situation and trends in structural changes and political conditions. In this phase, challenges and opportunities need to be identified.

The design phase is based on objectives defined for sustainable farming systems (Figure 1, phase 2a). These objectives are translated into five major themes: "quality production", "clean environment", "attractive landscape and diversified nature", "sustainable management of resources", and "farm continuity" (in this theme is next to farm economics also attention to labour and management aspects).

For each theme, a limited number of relevant parameters are defined on farm-level to characterise the systems performance. Each parameter is given a target value (Figure 1, phase 2b). The targets are future oriented and are derived from legislation, scientific evidence or expert knowledge. The themes together with the parameters and target values represent the objectives for the farming systems to be developed.

The next step is to design a suitable set of farming methods (Figure 1, phase 2c). Methods are defined here as coherent strategies on the major aspects of farming. Methods are for example "multifunctional crop rotation", "integrated nutrient management" and "integrated crop protection". In most cases, these methods need further development if they have to reach the objectives set.

The final step in the design phase is the development of a "theoretical prototype" combining all the selected methods (Figure 1, phase 2d). When completed, the prototype is checked, by expert judgement, comparing whether it is able to meet the targets set for the different parameters. The last part of this theoretical exercise shall provide detailed cropping programmes, allowing for adjustments that might be necessary for specific crops, weather and soil conditions.




Figure 1 Methodology of farming systems development in four phases. Phase 2 is divided in four sub-phases (after de Haan & Garcia Diaz, 2002)

The testing and improving phases are performed to verify the farming system in practice for at least a couple of years. Every year the actual value of the parameters has to be determined and compared to the target values. Shortfalls to the target values have to be analysed. Agronomic information and observations during the growing season are indispensable for this evaluation. The method(s) causing the shortfalls have to be identified and improved.

When, after testing and improving the system, the results are fulfilling (almost) all objectives, the results can be disseminated to all stakeholders involved in the production process (farmers, advisors, etc.) (Figure 1, phase 4). However, it is not the exact prototype to be disseminated, but the objectives with parameters and target values, the underlying integral approaches to farming, the farm methods and the toolbox with a practical guide to adjust the strategies to the local conditions.

Strong points of the prototyping method are the clear identification of shortfalls of the developed systems compared to the objectives set and identification of knowledge gaps. Weak points are the amount of data need and the time consuming process of testing and improving to reach a prototype that fits the objectives best.

Examples

1. *Redesign of farming systems because of insufficient solutions within current farming system. Example of the Netherlands*

In 2009 was concluded that in the Southeast of the Netherlands it is impossible to reach the EU Nitrates Directive and maintain yield and income of arable and vegetable farmers within the current farming systems (de Haan *et al.*, 2009; Meijer *et al.*, 2011). A redesign of systems has led to two new research pathways (de Haan *et al.*, 2010). The first is an agro-ecological pathway focused on improved soil management. This pathway shows that increased organic matter input can lead to increased yields and low nitrate leaching (de Haan & Verstegen, 2014). The second is a technical pathway, focused on soilless cropping. This pathway shows that soilless cropping of open field vegetables is possible and has perspectives for profitable production and large reduction in emissions (de Haan & van Dijk, 2013; Breukers *et al.*, 2014; de Haan *et al.*, 2014). This example shows that a shift of system perspective gives new opportunities, which are impossible within the current production system. However, still a lot of research and dissemination is necessary for a broad application of these new systems and strategies in practice.

2. *Two opposite approach of farming system are being carried out to solve fertiliser efficiency for leafy crops in open field conditions in Almeria*

The same shift in development of new farming systems is visible in Almeria (south-eastern Spain). Next to the conventional horticulture organic horticulture and hydroponic cultivation is in development (Rohner-Thielen, 2010). Research is done on both alternative farming systems. The hydroponic systems developed in Pulpi town (figure 2, left side) are based on exhaustive knowledge of mineral elements supply. The organic farming research executed in Nijar region (figure 2, right side) is based on organic matter management and its relationship with physical, chemical and biological soil conditions.

There are large differences in nutrient budgeting between the systems in terms of fertilisers (inputs), yield (outputs) and losses to the environment. There are some loss factors that do not exist in one of the extreme types of farming (organic and hydroponic), e.g. crop waste which is usually applied to the soil in organic horticulture or leaching which is limited in hydroponic because of the recycling of the nutrient solution. The efficiency of fertilisers in both organic and hydroponic system is higher compared to conventional farming, mainly because in organic farming no synthetic mineral fertiliser is used and nutrients are recycled continuously in closed hydroponics systems.

There is a large difference in control on nutrient uptake as well. In organic farming organic fertilisers are used and farmers trust on the release of nutrients for plant uptake through complex ecological interactions and biological processes (microorganism) during the growing season. In hydroponic systems there is more control possible on the composition of the nutrient solution. In conventional systems control is highly desirable, but difficult to realise with all complex soil processes. However various tools are in development to develop useful fertiliser programs.



Figure 2 Left side: Closed hydroponic system (NGS – New Growing System) in Pulpi (Almeria). Right side: organic horticulture in Níjar (Almeria).

3. Introduction of conservation agriculture in Hungary

The conservation agriculture medium-term programme started from 2014 with the participation of research units, seed and fertiliser companies and over 30 farms. The programme offers sustainable and intensive crop production solutions to increase the profitability of farmers. Sustainability and profitability requires crop cultivation techniques and economic management systems, to which the programme provides a solution with regard to all aspects, customised to the specific farm, by utilising the knowledge base of high tech companies. Preconditions in using the conservation agriculture practices (mostly natural and economic) are:

1. extreme weather conditions (drought or extremely wet year),
2. deteriorated soil conditions (soil structure and organic matter content),
3. increased production costs (cost of labour, fuel, input materials, etc.),
4. higher financing requirement for the more efficient and intensive cultivation techniques,
5. volatile commodity prices,
6. increased administrative tasks,
7. new criteria to participate in the single area subsidy payment scheme,
8. stricter environmental protection regulations.

The main elements of the integrated system are (Hadju, 2004a, b):

1. conservation tillage practices which are reducing climatic damage and improve soil physical structure, biological condition and water holding capacity, increase the nutrient supply capacity, better adapt the crop to the extreme weather conditions and reduce the cultivation cost,
2. optimising the nutrient supply according to the crop needs, with different intensity levels,
3. use of specific varieties and hybrids adapted to field conditions, emphasising early vigour and resistance to major diseases, adapted to specific nutrient management levels and to the changed tillage practices,
4. crop protection solutions adapted to the changed soil management practices (related to crop residues, weed control and moisture conditions) and nutrient management solutions which take into account the impacts of the previous crop, forecast-based applications, targeted use of pesticides (precision crop protection),

5. farm management software as an administrative basis for the efficient and transparent operation of the farm and supports the decision-making process based on local characteristics.

4. *Integrated production in Emilia-Romagna region in Italy*

In the last twenty years a system approach has been developed for several crops, including horticultural ones (fruits and vegetables), in the Emilia-Romagna region, in North Italy. The approach was established by the end of the '90s using the financial support of the Rural Development Plan, thus as a strategic tool in an effort of qualifying and increasing the quality of the agricultural production but respecting the environment. The system was developed by carrying on a detailed mapping of the soil chemical-physical properties of all land area of the Region, linking these data with the land use, soil capacity and water resources and producing zone maps (based on soil and weather characteristics, to identify suitable areas for different crops/cultivars). An extensive program of research about the crops nutrients needs, the test of varieties adapted to the different environments, and pest/diseases management based on Integrated Pest Management (IPM) was launched. The program was deeply linked with the activity of advisors and other professionals (e.g. the phytosanitary service of the Region) in order to promote the adoption of the different practices/measures that were aimed at reducing the overall environmental impact of the agricultural sector. The results of this strategic program can be assessed by considering that about 65% of the horticultural production of the region (more than 150.000 ha) is obtained by applying this integrated approach, which has led to the establishment of a quality system (QC – Certified Quality) by the Regional administration.

The farmers and advisors are supported in their decisions about the soil management (e.g. fertilisation, irrigation) by online applications that allow to calculate the amount of nutrients to be applied (e.g. <https://agri.regione.emilia-romagna.it/Suoli/>; <http://www.irriframe.it/>; Mannini *et al* 2013) on the basis of the crop history of the field, of the field physical-chemical characteristics, the precipitation history and weather forecasts, the expected yield of the crop, and the major pest/diseases of the specific crop (using for several of them also forecast models). The use of agronomical practices such as rotation, organic fertilisation (using also composts), fertigation, limited tillage, use of catch crops and alternative fertilisation strategies that are known to limit emissions and leaching of nutrients (e.g. mycorrhizal inocula, foliar fertilisation with organic fertilisers) are all recommended to be used within the different farming systems. The continuous search for new practices/products that can improve the production systems is provided to the farmers with new decision supporting tools which are also allowing for their economic evaluation.

Discussion

The inventories of measures in NUTRIHORT (Amery *et al.*, 2013) and Quemada *et al.* (2013) and Schoumans *et al.* (2014) provide a wide array of information about possible measures to increase nutrient use efficiency. Together with nutrient management measures, measures connected to other farming practices are summarised as well, e.g. measures on crop rotation, soil cultivation and water management. However, an inventory of measures aimed at one objective lacks by definition a systems approach. A system approach is by nature multi-objective and comprises multiple measures. Therefore, after the inventory, an evaluation of the feasibility of the different measures to a system needs to be made. To do this, elements from the prototyping approach may be used. Important aspects are the assessment of the current situation and the quantification of objectives of the farming systems.

The assessment of the current situation at farm and regional level should be based on: the main constraints, challenges and opportunities for the system, including biophysical (e.g. soil type and climate), socio-economic (e.g. farm structure and costs) and institutional (e.g. legislation and organisation of sales) aspects. From an agronomic point of view, an important aspect is related to the main growth limiting factors, which may have a larger effect on nutrient use efficiency than the application of a nutrient management measure (i.e. soil-borne diseases).

For a thorough evaluation of the effects of the measures on the farming system as a whole, it is needed to quantify the objectives of the farming system. This should include parameters and target values for all major themes within the farming system, not only those connected to the objective of nutrient efficiency to quantify side effects as well. If effects cannot be measured or estimated, research is needed to provide such data.

In the last few years, the attention to integrated soil management has increased strongly. It is expected that an integrated approach of soil chemical, physical and biological aspects will lead to a better performance of the farming system. In the farming systems development approach, integral soil management has still limited attention and methods connected to this subject have to be developed further, paying more attention to e.g. organic matter management and soil tillage.

For practical reasons of e.g. time and money, the choice can be made to not apply the whole prototyping method. This is visible in the examples presented. When using parts of the method large steps can be made efficiently. However, the risk is present that important aspects are overlooked. Most important is thus to think and act from a systems perspective.

Recommendations

From the prototyping methodology, the examples described above and discussions in the focus group, the following recommendations can be made to increase nutrient use efficiency from a systems perspective:

It needs to be made clear to farmers that a farm is a system and not a group of non-linked activities and that they have to analyse the whole system to get constraints, challenges and opportunities clear. When these are clear it will be easier to find solutions to challenges and opportunities. To strengthen the solutions the next issues should be encouraged to broaden the scope of the solutions:

- out of the box thinking,
- cooperation with other farmers and other stakeholders,
- emphasis on long term business and income stability.

The prototyping methodology does not necessarily have to be followed in all its details. It is important to follow the main steps in the methodology:

- Evaluate constraints, challenges and opportunities on biophysical, socio-economic and institutional aspects in the region of interest to assess the applicability of different measures.
- Estimate the most important growth limiting factors in the systems other than those related to fertilisation and when present improve these together with improving the fertiliser efficiency.
- Integrate measures in a farming system from a system perspective to assess all wanted and unwanted effects
- Define a parameter set with target values on all themes connected to the farming system and monitor the indicators.

To improve the prototyping methodology a method on integrated soil management should be developed.

Short term actions to improve systems thinking in vegetable growing can be the development of games from the (farming) models we have, e.g. get a realistic Hayday-game on horticulture and more interdisciplinary and international research and exchange of ideas and results.

References

- Amery, F., Vandecasteele, B., De Vis, R., Crappé, S., Van de Sande, T., Mechant, E., De Bolle, S., Willekens, K., De Neve, S., 2013. NUTRIHORT: Nutrient management, innovative techniques and nutrient legislation in intensive horticulture for an improved water quality. Fact sheets from the benchmark study on innovative techniques and strategies for reduction of nutrient losses in horticulture. September 16-18, 2013, Ghent, 130 p. [http://www.ilvo.vlaanderen.be/Portals/69/Documents/Book fact sheets NUTRIHORT.pdf](http://www.ilvo.vlaanderen.be/Portals/69/Documents/Book%20fact%20sheets%20NUTRIHORT.pdf)
- Breukers, M.L.H., Stokkers, R., Spruijt, J., Roelofs, P.F.M.M., Haan, J.J. de, 2014. Impact of deep flow cultivation systems on sustainability of field vegetable production. Acta Hort. (ISHS) 1034:169-176. http://www.actahort.org/books/1034/1034_20.htm
- Haan J.J., de (ed), 2002. Integrated and Ecological Nutrient Management. VEGINECO Report 3. Applied Plant Research. Lelystad. <http://edepot.wur.nl/11787>
- Haan J.J. de, Garcia Diaz, A. (eds), 2002. Manual on Prototyping Methodology and Multifunctional Crop Rotation. VEGINECO Report 2. Applied Plant Research. Lelystad. <http://edepot.wur.nl/11786>
- Haan, J.J. de, Zwart, K.B., Smit, A.B., Geel, W.C.A. van, 2009. Can intensive arable farming systems on sandy soils in the Netherlands meet the targets in the nitrate directive? Proceedings 16th Nitrogen Workshop, Connecting different scales of nitrogen use in agriculture, Turin, Italy, 28 June – 1 July 2009. pp. 471-472. <http://www.nitrogenworkshop2009.org/papers.htm>
- Haan, J.J. de, Sukkel, W., Stilma, E.S.C., 2010. Technology versus Agro-Ecology in Designing Vegetable Production Systems in the Netherlands. Acta Horticulturae 852. p. 123 - 130. <http://edepot.wur.nl/299920>
- Haan, J.J. de, Dijk, S.M. van, 2013. Soilless cultivation of outdoor horticultural crops in The Netherlands to reduce nitrogen emissions. In: Proceeding NUTRIHORT, 16-18 September 2013, Gent, België. - Gent, 2013 - p. 186 - 192. [http://www.vlm.be/SiteCollectionDocuments/Mestbank/Studies/Book_proceedings NUTRIHORT.pdf](http://www.vlm.be/SiteCollectionDocuments/Mestbank/Studies/Book_proceedings_NUTRIHORT.pdf)
- Haan, J.J. de, Spruijt, J., Vermeulen, T., 2014. Farm management and economic drivers for implementation of soilless cultivation of field vegetables in the Netherlands. Acta Hort. (ISHS) 1034:177-183. http://www.actahort.org/books/1034/1034_21.htm
- Haan, J.J. de, Verstegen, H., 2014. Effects of organic matter input on nitrate leaching and crop yield in arable and vegetable cropping on sandy soil in the Netherlands. In C.M.d.S. Cordovil. The Nitrogen challenge: building a blueprint for nitrogen use efficiency and food security. Proceedings 18th Nitrogen Workshop. Lisbon Portugal 30-6 – 3-7 2014. <http://www.nitrogenworkshop.com/#/abstracts/4585708037>
- Hajdu, Z., 2004a. Integrated crop production (booklet) ed. Soltub Bt. http://www.soltub.hu/?en_publications,31
- Hajdu, Z., 2004b. The application of the Good Agricultural Practices in vegetable production ed. (booklet) Soltub Bt. http://www.soltub.hu/?en_publications,31
- IFADATA, 2014. Production and International Trade. Paris: International Fertilizer Industry Association (IFA). Accessible via: <http://www.fertilizer.org/Statistics>. Accessed 04 September 2014.
- Mannini, P., Genovesi, R., Letterio, T., 2013. IRRINET: Large Scale DSS Application for On-farm Irrigation Scheduling. Procedia Environmental Sciences, 19: 823–829. DOI: 10.1016/j.proenv.2013.06.091.
- Meijer, B.J.M., Wijnands, F.G., Vogelesang, J.V.M., Stilma, E.S.C., Maas, M.P. van der, Jansma, J.E., Haan, J.J. de, Reuler, H. van, 2011. Systems innovations in agriculture in the Netherlands. Wageningen : WUR PPO-agv, 2011 (report PPO 340) - 51 p. <http://edepot.wur.nl/166572>
- Quemada, M., Baranski, M., Nobel-de Lange, M.N.J., Vallejo, A., Cooper, J.M., 2013. Meta-analysis of strategies to control nitrate leaching in irrigated agricultural systems and their effects on crop yield. Agriculture, Ecosystems and Environment 174, 1-10. DOI: 10.1016/j.agee.2013.04.018.
- Schoumans, O.F., Chardon, W.J., Bechmann, M.E., Gascuel-Oudou, C., Hofman, G., Kronvang, B., Rubaek, G.H., Ulén, B., Dorioz, J.-M., 2014. Mitigation options to reduce phosphorus losses from the agricultural sector and



improve surface water quality: A review. *Science of the Total Environment* 468-469, 1255-1266. DOI: 10.1016/j.scitotenv.2013.08.061.

Vereijken, P., 1999. Manual for prototyping integrated and ecological arable farming systems (I/EAFS) in interaction with pilot farms. Stichting Cereales. Wageningen. <http://edepot.wur.nl/17748>